

$$I_4^{\{D=4-2\epsilon\}}(0, m^2, p_3^2, m^2; s_{12}, s_{23}, 0, 0, m^2, m^2)$$

Page contributed by [R.K. Ellis](#)

Integral obtained from Eq.(A4), Equation 3 of ref. [1]

$$\begin{aligned} I_4^{\{D=4-2\epsilon\}}(0, m^2, p_3^2, m^2; s_{12}, s_{23}, 0, 0, m^2, m^2) &= \left(\frac{\mu^2}{m^2}\right)^\epsilon \frac{1}{(s_{12} - m^2)(s_{23} - m^2)} \\ &\times \left[ \frac{1}{\epsilon^2} - \frac{1}{\epsilon} \left\{ \ln\left(1 - \frac{s_{23}}{m^2}\right) + \ln\left(1 - \frac{s_{12}}{m^2}\right) \right\} - \ln^2(x_{p_3^2}) \right. \\ &\quad \left. + 2 \ln\left(1 - \frac{s_{12}}{m^2}\right) \ln\left(1 - \frac{s_{23}}{m^2}\right) - \frac{\pi^2}{2} \right] + \mathcal{O}(\epsilon) \end{aligned}$$

See the file on [notation](#). We make use of the quantities

$$\beta_r = \sqrt{1 - \frac{4m^2}{r}}, \quad x_r = \frac{\beta_r - 1}{\beta_r + 1},$$

where  $r$  represents a generic invariant. Analytic continuation by the replacements  $s_{ij} \rightarrow s_{ij} + i\epsilon, p_j^2 \rightarrow p_j^2 + i\epsilon$ .

See also ref. [2], Eq. (30)

[Return to general page on boxes](#)

## References

- [1] W. Beenakker, S. Dittmaier, M. Kramer, B. Plumper, M. Spira and P. M. Zerwas, Nucl. Phys. B **653**, 151 (2003) [[arXiv:hep-ph/0211352](#)]. [[arXiv:hep-ph/0211352](#)]
- [2] G. Rodrigo, A. Santamaria and M. S. Bilenky, J. Phys. G **25**, 1593 (1999) [[arXiv:hep-ph/9703360](#)]