

$$I_4^{\{D=4-2\epsilon\}}(0, p^2, 0, p^2; t, u; 0, 0, m^2, m^2)$$

Page contributed by [R.K. Ellis](#)

This integral has been given in Eq. (6.72) of ref [1]. In the region $t, u, p^2 < 0$ and $m^2 > 0$ we have,

$$I_4^{\{D=4-2\epsilon\}}(0, p^2, 0, p^2; t, u; 0, 0, m^2, m^2) = \frac{1}{[(m^2 - t)(m^2 - u) - (m^2 - p^2)^2]} \left[\left[\frac{1}{\epsilon} - \ln \left(\frac{(m^2 - p^2)^2}{\mu^2 m^2} \right) \right] \ln \left(\frac{(m^2 - p^2)^2}{(m^2 - t)(m^2 - u)} \right) - \ln^2 \left(\frac{m^2 - p^2}{m^2 - t} \right) - \ln^2 \left(\frac{m^2 - p^2}{m^2 - u} \right) - 4 \operatorname{Li}_2 \left(1 - \frac{m^2 - p^2}{m^2 - t} \right) - 4 \operatorname{Li}_2 \left(1 - \frac{m^2 - p^2}{m^2 - u} \right) - 2 \operatorname{Li}_2 \left(1 - \frac{(m^2 - t)(m^2 - u)}{(m^2 - p^2)^2} \right) \right] + \mathcal{O}(\epsilon)$$

For Li_2 etc, see the file on [notation](#).

[Return to general page on boxes](#)

References

- [1] R. Höpker, Hadroproduction and decay of squarks and gluinos, (in german), DESY Internal report DESY-T-96-02, ([Relevant excerpt](#))