

$$I_3^{\{D=4-2\epsilon\}}(0, 0, p^2; 0, 0, 0)$$

Page contributed by [R.K. Ellis](#)

Page verified by [Lance Dixon](#)

Triangle with two massless external lines

$$\begin{aligned} I_3^{\{D=4-2\epsilon\}}(0, 0, p^2; 0, 0, 0) &= \frac{\mu^{2\epsilon}}{\epsilon^2} \left( \frac{(-p^2 - i\epsilon)^{-\epsilon}}{p^2} \right) \\ &= \frac{1}{p^2} \left( \frac{1}{\epsilon^2} + \frac{1}{\epsilon} \ln \left( \frac{\mu^2}{-p^2 - i\epsilon} \right) + \frac{1}{2} \ln^2 \left( \frac{\mu^2}{-p^2 - i\epsilon} \right) \right) + \mathcal{O}(\epsilon). \end{aligned}$$

For  $\epsilon, \epsilon$  see the file on [notation](#).

We also present this function with the singularities regulated with a small mass  $\lambda$ , (from ref. [\[1\]](#), Eq.(A.3))

$$I_3^{\{D=4\}}(0, 0, p^2; \lambda^2, \lambda^2, \lambda^2) = \frac{1}{2p^2} \ln^2 \left( \frac{-p^2 - i\epsilon}{\lambda^2} \right) + \mathcal{O}(\lambda)$$

[Return to general page on triangles](#)

## References

- [1] W. Beenakker, S. Dittmaier, M. Kramer, B. Plumper, M. Spira and P. M. Zerwas, Nucl. Phys. B **653**, 151 (2003) [[arXiv:hep-ph/0211352](#)]. [[arXiv:hep-ph/0211352](#)]